

Specimen Label



®Trademark of Dow AgroSciences LLC

A selective herbicide for the control of broadleaf weeds in field corn.

Active Ingredients:

flumetsulam: <i>N</i> -(2,6-difluorophenyl)-5- methyl-1,2,4-triazolo-[1,5a]- pyrimidine-2-sulfonamide	18.5%
clopyralid potassium salt: 3,6-dichloro-2-pyridinecarboxylic acid, potassium salt	60.0%
Other Ingredients	21.5%
Total Ingredients	100.0%

Acid Equivalent: clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid – 50%

Contains 0.185 pound flumetsulam active ingredient and 0.5 pound clopyralid acid equivalent per pound of product.

EPA Reg. No. 62719-315

Keep Out of Reach of Children

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled, Or Absorbed Through The Skin • May Cause Skin Sensitization Reactions In Certain Individuals

Do not get in eyes or on clothing. Avoid breathing vapors or spray mist and contact with skin.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA Chemical Resistance Category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

The active ingredients in this product are known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this product where soils are permeable, particularly where the water table is shallow, may result in leaching to ground water.

Caution should be exercised when handling this product at mixing and loading sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read “Warranty Disclaimer,” “Inherent Risks of Use,” and “Limitation of Remedies” elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

NOT FOR SALE, USE OR DISTRIBUTION IN NASSAU AND SUFFOLK COUNTIES IN NEW YORK STATE.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 122°F for extended periods of time. If container is damaged or spill occurs, use product immediately or contain with absorbent materials and dispose as waste. **Water Soluble Packaging:** Packets may become brittle when stored below 32°F. Handle carefully when frozen to avoid breakage or allow package to warm above 32°F before handling.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site according to label use directions or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable nonrigid containers:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Refillable rigid containers larger than 5 gal:

Container Reuse: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Storage and Disposal (Cont.)

Nonrefillable rigid containers larger than 5 gal:

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times.

Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specific minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Do not apply this product through any type of irrigation system.

Do not use flood irrigation to apply or incorporate this product.

Product must be used in a manner that will prevent back-siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

General Information

Hornet® WDG broadleaf blend herbicide is a selective herbicide for broadleaf weed control in field corn, including high oil and waxy varieties and field corn grown for ensilage. Hornet WDG may be applied as a preplant surface, preplant incorporated, preemergence, or postemergence treatment. Soil surface treatments may be applied with water, liquid fertilizer, or impregnated on dry fertilizer. Postemergence treatments should be applied with water. Absorption of Hornet WDG occurs from both shoot and root uptake. Susceptible weeds exposed to Hornet WDG stop growing and either die or remain non-competitive with the crop. Hornet WDG provides residual control of weeds that may emerge after application. Adequate soil moisture is necessary for optimal activation because uptake and translocation of Hornet WDG involves uptake by emerging shoots and/or roots.

Use directions in Dow AgroSciences supplemental labeling may supersede directions or limitations in this labeling.

Use Precautions and Restrictions

Handling Precautions if Product is Packaged in Water Soluble

Packets: Do not remove water soluble packet from overpack except for immediate use. Do not allow water soluble packet to come into contact with water prior to use. Do not handle water soluble packet with wet hands or wet gloves. Carefully reseal package containing unopened water soluble packets and protect package from moisture.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Application Precautions

- Uneven application or uneven incorporation of Hornet WDG can result in erratic weed control or crop injury. Over application may result in crop injury or rotational crop damage from soil residue.

Adverse Weather Conditions

- **Soil Application Only:** Extended cold, wet conditions (soil temperatures below 50°F and excessive rainfall with wet soil conditions), following soil application of Hornet WDG to field corn, which persist during germination and/or early crop development may result in crop injury. Injury symptoms, which include yellowing of leaves and/or crop stunting, are usually temporary and affected corn plants usually recover without affecting yield.
- When applications are made under adverse (dry or cold) conditions or when large weeds or less susceptible species are treated, only weed suppression may be observed. Weed suppression is a visual reduction in weed competition (reduced population, size, and/or vigor) as compared to an untreated area. Degree of control can be increased by applying Hornet WDG under favorable growing conditions (i.e., adequate moisture and temperature), and by using a higher rate in the specified rate range.
- Dry weather following preplant surface or preemergence applications of Hornet WDG may reduce effectiveness. If sufficient activating rainfall or overhead irrigation does not occur within 7 to 10 days of application, rotary hoe, harrow, or shallowly cultivate to incorporate the herbicide lightly into the soil. Use a preplant incorporated application when a period of dry weather is predicted after application.
- Avoid application when air temperature is near freezing or when freezing conditions are expected for several days following application.
- Postemergence application of Hornet WDG to corn that is stressed or damaged by conditions such as cold weather, hot weather (>90°F), hail, drought, water saturated soil, disease, or insects may cause crop injury.

Tank Mixing

Hornet WDG may be tank mixed or followed by other overlay or postemergence treatments registered for use on corn to broaden the spectrum of weeds controlled. This product may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing is not prohibited by the label of the tank mix product; and (3) the tank mix combination is compatible as determined by a "jar test" described in the "Tank Mix Compatibility Testing" section below.

Tank Mixing Precautions:

- Read carefully and follow all applicable use directions, precautions, and limitations on the respective product labels.
- Do not tank mix Hornet WDG with Basagran herbicide, Laddock herbicide, or Lightning herbicide as severe crop injury may occur. (See instructions for Postemergence Treatments, Tank Mixing.)
- Do not exceed recommended application rates. Do not tank mix with another pesticide product that contains the same active ingredient as this product unless the label of either tank mix partner specifies the maximum dosages that may be used.
- For products packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a product mixture containing boron unless the tank and spray equipment has been adequately cleaned. (See instructions for Sprayer Clean-Out.)

Tank Mix Compatibility Testing: A jar test is recommended prior to tank mixing to ensure compatibility of Hornet WDG and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, jels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Restrictions and Precautions for Soil Applications of Hornet WDG (Not Applicable to Postemergence Use)

- **Corn Planting Depth:** Minimum planting depth should be at least 1 1/2 inches.
- Do not soil apply to peat or muck soils as reduced weed control will result. (May be used postemergence.)
- Do not apply to areas where the soil pH is greater than 7.8 as this may result in increased crop injury.
- Do not apply to a soil containing greater than 5% organic matter if the soil pH is below 5.9 as reduced weed control will result.
- Use of Hornet WDG in soil-applied treatments on soils with less than 1.5% organic matter (O.M.) may result in crop injury. Apply as a soil-treatment to fields which have less than 1.5% O.M. only if the risk of crop injury is acceptable.
- If any herbicide with ALS (acetolactate synthase) inhibition mode of action such as Pursuit, Preview, Canopy, Classic, Scepter, or Squadron herbicide, etc., was applied the previous year, apply Hornet WDG to corn only if the rotational restrictions applicable to corn for the preceding product has been met.

- Corn growing in calcareous soils or soils with historically high salt content (soil test results for salinity indicating electrical conductivity greater than 1.0 mmho/cm) may exhibit chlorosis and/or stunting resulting from reduced availability of iron, zinc or other micro nutrients essential for normal crop vigor and growth. The presence of soil-active herbicides, such as Hornet WDG may cause additional stress under these conditions resulting in increased leaf chlorosis and/or crop stunting. This added stress may retard crop recovery, especially under conditions of limited rainfall. In fields, which contain calcareous or high salt content soils, growers should plant "IR" or IMR" designated varieties, commonly referred to as "imidazolinone resistant" corn hybrids. On these type soils, the likelihood of crop injury can also be reduced by using the lower end of the recommended rate range for the soil type and/or by applying Hornet WDG 10-14 days prior to planting.

Soil Insecticide Advisories for Soil Applications of Hornet WDG:

When Hornet WDG is used for soil applied broadleaf weed control in corn:

- Soil applied organophosphate insecticides should be applied in a T-band or a band to avoid potential crop injury.
- Soil insecticides from other classes of chemistry may be applied in-furrow, T-banded, or banded.
- Terbufos (Counter insecticide products) or phorate (Thimet insecticide products) should not be used.

Soil Insecticide Advisories for Postemergence Applications of Hornet WDG:

- Do not apply Hornet WDG postemergence if corn was previously treated with Counter insecticide or Thimet insecticide as severe crop injury may result.
- Postemergence applications of Hornet WDG to corn previously treated with T-band, band, or in-furrow applications of other organophosphate insecticides such as Lorsban® insecticide, Aztec, Fortress, or Dyfonate insecticides may cause temporary crop injury.

Foliar Insecticide Advisories for Postemergence Applications of Hornet WDG

- Do not tank mix Hornet WDG with foliar postemergence organophosphate insecticides as severe crop injury may result. To avoid crop injury, apply the foliar organophosphate insecticide treatment at least 10 days before or 10 days after the application of Hornet WDG.
- Hornet WDG may be tank mixed with non-organophosphate foliar insecticides provided they are labeled for use with postemergence corn herbicides.

Use with other Products

- Corn previously treated with Hornet WDG that is stressed or damaged by conditions such as cold weather, hail, drought, water saturated soil, disease, or insects should not be treated with Accent, Beacon, Permit, Exceed, or Basis herbicides, or other herbicides with ALS inhibition mode of action as this may cause further crop injury.
- Do not foliar apply Hornet WDG to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

Use with Genetically Modified Corn Varieties

- If an "IR" or "IMR" designated hybrid (commonly referred to as "imidazolinone resistant") is planted, any organophosphate insecticide, including Counter or Thimet, can be applied according to label directions without increasing the likelihood of injury to corn from Hornet WDG. The adverse interaction between Counter or Thimet insecticide and Hornet WDG **does not** occur in corn hybrids identified as "IR" or "IMR". This adverse interaction **does** occur in imidazolinone tolerant "IT", "PT" hybrids which are considered as "standard" hybrids regarding this effect. "IR" or "IMR" hybrids may also be planted to reduce injury to corn from preemergence treatments of Hornet WDG on soils with less than 1.5% organic matter or pH greater than 7.8.

Maximum Application Rate

- Do not exceed a total application rate of 6.0 oz per acre of Hornet WDG (0.07 lb a.i. of flumetsulam) in a single crop year.
- Multiple applications of Hornet WDG within a growing season can be made as a soil application followed by a postemergence application, or as multiple postemergence applications. Do not exceed the cumulative rate of 0.07 lb per acre active ingredient of flumetsulam per single crop year if a postemergence application of Hornet WDG is made following a soil application of a flumetsulam-containing herbicide or with a postemergence herbicide containing flumetsulam (See table below to calculate cumulative flumetsulam amount per season.).
- Do not exceed a cumulative amount of 0.25 lb a.i. per acre of clopyralid per single crop year. (See table below to calculate cumulative clopyralid amount per season.)

Examples: 4.0 ounces of Hornet WDG contains 0.047 lb flumetsulam and 0.125 lb clopyralid.
 2.9 ounces of Accent Gold herbicide contains 0.035 lb flumetsulam and 0.094 lb clopyralid.

Herbicide	Unit of Measure	Flumetsulam (lb a.i./ unit of measure)	Clopyralid (Acid) (lb a.i./unit of measure)
Hornet	1 ounce	0.0145	0.039
Hornet WDG	1 ounce	0.0116	0.031
Python® WDG	1 ounce	0.05	---
Accent Gold	1 ounce	0.012	0.032
Stinger®	1 fluid ounce	---	0.023

The maximum active ingredient allowed per season in all states:
 Flumetsulam = 0.07 lb/acre
 Clopyralid = 0.25 lb/acre

NOT FOR SALE, USE OR DISTRIBUTION IN NASSAU AND SUFFOLK COUNTIES IN NEW YORK STATE.

Other Precautions and Restrictions

- Do not apply Hornet WDG to sweet corn or popcorn.
- **Hybrid Seed Production:** Corn inbred lines grown for hybrid seed production may be injured by Hornet WDG. Inbred lines should be thoroughly tested for crop tolerance before treating large acreage. While growers are not prohibited from using Hornet WDG on seed corn, **Dow AgroSciences will not accept responsibility for any crop injury arising from the use of Hornet WDG on field corn grown for seed.**
- **Preharvest interval:** An interval of at least 85 days is required between application of Hornet WDG and field corn harvested for grain. If field corn is grown for forage or ensilage, application must occur before corn reaches 20 inches in height or V6 growth stage (whichever occurs first) and an interval of at least 45 days is required between application and harvest.

- **Do not aerially apply Hornet WDG unless permitted by EPA approved supplemental labeling.**
- **Avoid all direct or indirect contact with nontarget plants.** Do not apply near desirable vegetation. Allow adequate distance between target area and desirable plants under conditions of application to minimize potential exposure.
- **Crop Residues from Treated Areas:** Crop residues from treated areas cannot be used for composting or mulching on ground where susceptible crops may be grown the following season. To promote herbicide decomposition, plant material should be evenly incorporated or burned. Adequate moisture is also required to promote breakdown of plant residues, which contain clopyralid.
- **Do not move treated soil.** Avoid situations where soil particles may blow into areas where susceptible crops are grown. The hazard of movement of this product on dust is reduced if treated fields are irrigated or if rain occurs shortly after application.
- **Do not apply under conditions that favor runoff or wind erosion of soil containing Hornet WDG to nontarget areas. To prevent off-site movement due to runoff or wind erosion:**
 - Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
 - Do not apply to impervious substrates such as paved or highly compacted surfaces or frozen or snow covered ground.
 - Do not apply to soils when saturated with water.
 - Do not use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.
- **Do not apply when weather conditions favor drift to nontarget sites.** Spray drift of Hornet WDG to emerged soybeans or soil to which soybeans will be planted during the same growing season may cause soybean injury.
- **Read and follow these Advisories to minimize drift to nontarget areas.**
 - Minimize drift by using sufficient spray volume to ensure adequate coverage with large-droplet size sprays.
 - Use low pressure application equipment capable of producing a large-droplet spray. Do not use nozzles that produce a fine-droplet spray. Droplet size has been shown to be the single most important factor affecting drift from ground applications.
 - While increasing droplet size does reduce the potential for spray drift, larger droplets do not eliminate drift if environmental or application conditions are inappropriate for application.
 - Use larger capacity nozzles to increase flow rate rather than increasing spray pressure.
 - Keep height of ground-driven spray booms as low as possible above the target to minimize exposure to evaporation and wind while still providing good coverage. Applications made late in the growing season with excessive boom heights drastically increase the potential for spray drift.
 - Do not apply when wind is gusting or wind speed exceeds 15 mph as uneven spray coverage and drift may result. Avoid application to border rows adjacent to susceptible crops such as soybeans, field peas, or sunflowers under windy conditions unless one of the following drift management steps is taken:
 - (1) application is made only when the wind direction is such that the susceptible crop is up-wind from the treatment area (wind blowing from the susceptible crop toward the treated crop); or
 - (2) the applicator leaves an adequate buffer zone between the treated crop and the susceptible crop and coarse or low drift nozzle configurations are used.

A drift control or deposition agent may be used with this product to aid in reducing spray drift due to wind when making applications adjacent to susceptible crops, but may not be effective after prolonged pumping of the spray mix.

- On calm days with little or no wind, check for temperature inversions before making herbicide applications. Temperature inversions occur under calm conditions with little or no wind and air temperature increases with increasing height above the ground. Inversion conditions may be indicated by a layer of fog or mist near the ground and, under clear conditions, may be detected by use of a smoke column. A temperature inversion is indicated when smoke does not rise in a column, but layers at some level above the ground. Do not apply herbicides if temperature inversion conditions exist in the treatment area.

Sprayer Cleanup

To avoid injury to or exposure of nontarget crops, thoroughly clean and drain spray equipment used to apply Hornet WDG after use. Cleaning should occur as soon as possible after application of Hornet WDG. Spray equipment should be cleaned after use with Hornet WDG by the following procedure:

1. Drain any remaining Hornet WDG from the spray tank and dispose of according to label disposal instructions.
2. Hose down the interior surfaces of the tank. Flush tank, hoses, boom, and nozzles with clean water for 10 minutes. Fill the tank with water and recirculate for 15 minutes. Spray part of the mixture through the hoses, boom, and nozzles and drain the tank. All rinse water must be disposed of in compliance with local, state, and federal guidelines.
3. Fill the tank with water and recirculate for 15 minutes. For optimum cleaning, a tank cleaner such as liquid ammonia (1 gallon per 100 gallons of water) or other commercial tank cleaner is recommended in the second rinse if the spray equipment will be used on crops other than field corn. Spray part of the mixture through the hoses, boom, and nozzles and drain the tank. All rinse water must be disposed of in compliance with local, state, and federal guidelines.
4. Remove the nozzles and screens and clean separately.
5. If the spray equipment will be used on crops other than field corn, repeat steps 1 and 2 again and thoroughly wash the spray mixture from the outside of spray tank and the boom.

Rotational Crop Restrictions

When tank mixing with companion herbicides, follow the most restrictive crop rotation guidelines on the label of each product used.

The following rotational crops may be planted after the indicated interval following application of rates up to 6.0 ounces per acre of Hornet WDG:

Numbers within parentheses (-) in table refer to Specific Rotational Crop Requirements below.

Rotational Crop	Interval (Months)
barley, oats, rye, wheat	4
alfalfa (1), dry beans (1), forage grasses (2), lima beans (1), popcorn, rice, seeding of cover crops (3), soybean (1)	10.5
grain sorghum	12
peas (1, 4), snap beans (1, 4)	18
cotton, peanuts, potatoes, sunflower, sweet corn (5), tobacco	18
sugar beets, canola and all other crops (6)	26

Specific Rotational Crop Requirements:

1. When annual rainfall and/or irrigation is less than 15 inches on soils with less than 2% organic matter, alfalfa, dry beans, lima beans, peas, snap beans, and soybeans should not be planted until 18 months after treatment.
2. Excludes forage grasses grown for commercial seed production.
3. The following cover crops may be planted for establishment of Federal Conservation Reserve Programs and Agricultural Reserve Programs no sooner than 10.5 months following application of Hornet WDG at rates up to 4.0 oz per acre: **legumes** including alfalfa, clovers, crownvetch, birdsfoot trefoil, and lespedeza; and **grasses**, including big bluestem, little bluestem, switchgrass, Russian wildrye, green needle, smooth bromegrass, Garrison creeping foxtail, canary grass, orchardgrass, intermediate wheatgrass, tall wheatgrass, crested wheatgrass, western wheatgrass and indian grass.
Some stand reduction or temporary stunting of legume seedlings is possible. However, **Dow AgroSciences will not** accept responsibility for any crop injury or stand failure in crops established under Federal Conservation Reserve Programs and Agricultural Reserve Programs following use in corn and the subsequent 10.5 month rotational crop restriction. Additionally, Dow AgroSciences will not accept responsibility for any crop injury or stand failure of native grasses as a result of inadequate seedbed preparation, erratic germination, lack of seedling vigor, or plant stress from unfavorable environmental conditions.
4. An 18-month crop rotation is recommended following application of Hornet WDG at rates **greater** than 4 oz per acre. Peas and snap beans may be planted 10.5 months following application of Hornet WDG at rates up to 4 oz per acre.
5. **Certain sweet corn varieties** may be planted 10.5 months following soil or postemergence application of up to 4.0 oz per acre of Hornet WDG. This interval applies only to the following varieties of sweet corn grown for processing: Bonus, Challenger, Chase, Cornucopia, Crisp'N Sweet 710, Crisp'N Sweet 710A, DMC 20-04, DMC 20-35, Eliminator, Empire, Excalibur, Excellency, GH 0937, GH 2547, GH 2628, GH 2690, GG 5, GG 8, GG 22, GG 23, GG 40, GG 43, GG 46, GG 55, GG 246, GG 255, GG 256, GG 539, HM 701, 781 Ultra, Lumina, Reward, Sheba, Spirit, Sprint, Viking, and Zenith. The rotational interval is 18 months for other sweet corn varieties not listed here, except as provided in updated listings of sweet corn varieties tolerant to this product.
6. Rotation to sugar beets, canola, and all other crops requires a 26-month rotation interval and a successful field bioassay.

Field Bioassay Instructions: In fields previously treated with this product, plant short test rows of the intended rotational crop across the original direction of application in a manner to sample field conditions such as soil texture, soil pH, drainage, and any other variable that could affect the seed bed of the new crop. Field bioassay at any time between harvest of the treated crop and the planting of the rotational crop. Observe the test crop for herbicidal activity, such as poor stand (effect on seed germination) chlorosis (yellowing), and necrosis (dead leaves or shoots), or stunting (reduced growth). If herbicidal symptoms do not occur, the test crop can be grown. If there is apparent herbicidal activity, do not plant the field to the test rotational crop; plant only a labeled crop or crop listed in the table above for which the rotational interval has clearly been met.

Mixing and Application

Spray Volume

Apply Hornet WDG in sufficient spray volume to provide uniform coverage using properly calibrated ground equipment. Apply in a total spray volume of 10 to 60 gallons per acre using low pressure (20-40 lb/sq in). Maintain sufficient agitation during mixing and spraying to ensure a uniform spray mixture. More thorough coverage is possible when making soil applications to minimum or no-till corn by using a total spray volume of 20 or more gallons per acre.

Hornet WDG (oz/acre)	Acres per Package	
	Acres Per 6 lb Plastic Jug †	Acres per 6 oz Packet (Contains four 1.5 oz units) ††
2.0	48	3.0
3.0	32	2.0
4.0	24	1.5
5.0	19.2	1.2

† If the number of acres to be treated results in the use of a partial container, use the measuring device provided with the container to measure out product according to the scale indicated on the measuring device.

†† To calculate the number of 6 oz water soluble packets for your spray mix:

- Determine the number of acres you wish to spray in the desired application.
- Divide the number of acres by the acres per packet that falls within the rate range you have chosen. See the above table for broadcast application rates and corresponding acres per packet.
- The result is the number of packets you are required to add to the spray mix.

If the resulting number of packets is not a whole packet:

- Do not open the water soluble packets.**
- Round up or down to the nearest whole number of packets and check to make sure that the resulting number of acres per packet falls within the desired rate range for the application.

Sample Calculations:

- Planned application = 3.0 oz per acre (The acres per packet = 2.0).
- Assuming 17 acres is to be treated, 17 acres divided by 2.0 acres per packet = 8.5 packets (Round up to 9 packets).

Band Application

Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre} = \text{Amount needed per acre of field}$$

Mixing Directions

Hornet WDG is a water dispersible granule formulation. Thorough mixing is required.

- Fill the tank with 1/2 of the total amount of water or liquid fertilizer required for the load.
- Start agitation system.
- Add the required amount of Hornet WDG directly into the spray tank while agitating. If product is packaged in water soluble packets, open the overpack and add the required number of water soluble packets directly to the spray tank while agitating. **(For use of water soluble packaging in liquid fertilizer solutions, see the "Application in Liquid Fertilizer" section of this label for special pre-mixing instructions.)** Do not open water soluble packets. Water soluble packets will float on the surface until the water soluble film dissolves and releases the product. Handling packets with hands should be minimized.
- Continue agitation and complete filling the tank while product disperses in the spray tank solution.

Before spraying make sure Hornet WDG is thoroughly mixed in the solution. If product is in water soluble packets, make sure packets have completely disintegrated and product is thoroughly mixed with water. Depending on the water temperature and the degree of agitation, the packet and Hornet WDG should be completely dispersed within 5 minutes from the time they were added to the water.

To insure a uniform spray mixture continuous agitation is required during mixing and spraying. Apply within 24 hours after mixing. If product is allowed to settle, thoroughly agitate to resuspend the mixture before spraying.

Hornet WDG in Tank Mix

Hornet WDG may be applied in tank mix combination with labeled rates of other products provided (1) the tank mix product is labeled for the timing and method of application for the use site to be treated; and (2) tank mixing with Hornet WDG is not prohibited by the label of the tank mix product. See "Tank Mixing" in "General Use Precautions" section.

Vigorous, continuous agitation during mixing, filling, and throughout application is required for all tank mixes. Sparger pipe agitators generally provide the most effective agitation in spray tanks. To prevent foaming in the spray tank, avoid stirring or splashing air into the spray mixture.

Mixing Order for Tank Mixes: Fill the spray tank to 1/4 to 1/3 of the total spray volume required with water or liquid fertilizer solution. Start agitation. Add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: Hornet WDG (slurried if mixing water soluble packets with liquid fertilizer) and other dry flowables; wettable powders; aqueous suspensions, and flowables. Maintain agitation and fill spray tank to 3/4 of total spray volume. Then add emulsifiable concentrates and any solutions.

Note: Spray adjuvants (non-ionic surfactants, crop oil concentrates, methylated seed oil, urea ammonium nitrate, and ammonium sulfate) required for postemergence foliar applications should be added to the spray tank last.

Finish filling the spray tank. Maintain continuous agitation during mixing, final filling, and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled material may be more difficult to resuspend than when originally mixed.

Line screens in the spray tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

Application with Liquid Fertilizer

Bottled Hornet WDG is the preferred product form for use in liquid fertilizer. Hornet WDG in water soluble packets can be used, but **must be premixed or slurried with water prior to use in liquid fertilizer**. It is important that **all water soluble packet material be totally dissolved before transferring into liquid fertilizer**. Any packet material undissolved in the premix will remain undissolved in the liquid fertilizer solution and could potentially lead to the clogging of screens and nozzles. For best results, use a minimum of 2 pints of water for every 6 oz of Hornet WDG water soluble packets. Add the Hornet WDG packets while mixing and **continue mixing until all packet material is dissolved** and granules are dispersed. The time needed to completely dissolve the packet material will depend upon water temperature and efficiency of mixing. Increasing the amount of premix water will decrease required mixing time. It is also recommended that premix and rinsate from the premix container be added to the spray tank through a 20–35 mesh screen.

When necessary, a compatibility agent can be used to ensure that Hornet WDG mixes properly. The use of appropriate compatibility agents is especially important when tank mixing Hornet WDG and other dry flowables, wettable powders, flowables, liquids, aqueous suspensions, or solutions with emulsifiable concentrates in liquid fertilizers. If the emulsifiable concentrate formulation rises to the surface of the fertilizer as an oil (“oils out”), the oil may combine with the wettable powder, flowable, or suspension to form oily curds (viscous phase) which are difficult to disperse. A jar test, utilizing relative proportions of the tank mix ingredients is recommended prior to mixing with liquid fertilizers.

Note: Do not use liquid fertilizer as the carrier when Hornet WDG is applied postemergence to corn.

Application with Dry Bulk Fertilizer

Dry bulk fertilizer may be impregnated or coated with Hornet WDG. Application of dry bulk fertilizer impregnated with Hornet WDG provides weed control equal to the same rates of Hornet WDG applied in liquid carriers. Follow label recommendations for Hornet WDG regarding rates per acre, crops, special instructions, cautions, and special precautions. Apply 200 to 700 pounds of the fertilizer/herbicide mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Non-uniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control.

Most dry fertilizers can be used for herbicide impregnation with Hornet WDG. When coated ammonium nitrate and/or limestone are used alone, do not impregnate with Hornet WDG; these materials will not absorb the herbicide. Fertilizer blends containing coated ammonium nitrate and/or limestone as a part of the fertilizer mixture can be impregnated.

Compliance with all federal and state regulations relating to blending pesticide mixtures with dry bulk fertilizer, registration, labeling, and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale.

Impregnation: Hornet WDG must be pre-mixed or slurried with water prior to impregnation of dry bulk fertilizer. For best results, use a minimum of 2 pints of water per 6.0 oz water soluble packet. To make the water slurry, add the required rate of Hornet WDG (see formula below) to enough water to give a total volume of at least 6 pints of solution per ton of fertilizer. Make sure the Hornet WDG is thoroughly dispersed in the water before spraying onto the fertilizer. Spray nozzles should be placed to provide uniform spray coverage onto the fertilizer. Care should be taken to aim the spray directly onto the fertilizer and avoid spraying the walls of the blender. Use any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender.

Calculate amounts of Hornet WDG by the following formula:

$$\frac{2,000}{\text{Pounds/acre of fertilizer}} \times \frac{\text{Pounds/acre of Hornet}}{\text{of Hornet}} = \frac{\text{Pounds of product}}{\text{per ton of fertilizer}}$$

Note: Thoroughly clean dry fertilizer blending equipment prior to use with other herbicides. It is important to clean the blender, herbicide spray tank, and spraying apparatus thoroughly. Rinse the sides of the blender and the herbicide tank with water. Then, impregnate the rinsate onto a load of dry fertilizer intended for an approved crop. Use a maximum rate of 1 gallon of rinsate per ton of fertilizer. Follow with 1 to 2 loads of unimpregnated fertilizer in the blender before switching herbicides. The fertilizer application equipment must be empty, clean, and dry before applying any material to crops other than corn.

Crop-Specific Use Directions

FIELD CORN

Soil Applied Treatments

Broadcast Application Rates (Preplant Surface Applied, Preplant Incorporated, Postplant Preemergence, and Spike Stage Treatments)

Soil Texture	Hornet WDG (oz/acre)	
	<3.0% Organic matter	>3.0% Organic matter
Coarse	4.0	4.0 - 5.0
Medium or Fine	4.0 - 5.0	5.0

Note: Use the high end of the rate range on soils with greater than 3% organic matter and/or when applications are made 14 to 30 days before planting.

Broadleaf Weeds Controlled by Hornet WDG when Soil Applied
Hornet WDG will control “triazine tolerant” biotypes of these weeds, commonly know as “triazine resistant”.

Note: Numbers within parentheses (-) in weeds list refer to “Use Information for Specific Weeds” below.

amaranth, Palmer	pigweed, smooth
anoda, spurred	poinsettia, wild
beggarweed, Florida	puncturevine
buckwheat, wild	purslane, common
carpetweed	ragweed, common
chickweed, common	ragweed, giant(1)
cocklebur, common	shepherd’s purse
clover, red	sicklepod
henbit	sida, prickly
horseweed (marestail)	smartweed, Pennsylvania
jimsonweed	spurge, nodding
kochia(1, 5)	spurge, prostrate
ladythumb	spurge, spotted
lambsquarters, common	sunflower, common
mallow, Venice	thistle, Canada (3)
morningglory, entireleaf (1)	velvetleaf
morningglory, ivyleaf (1)	waterhemp species (4, 5)
morningglory, tall (1)	wormwood, biennial
mustard, wild	
nightshade species(2)	
pigweed, redroot	

Use Information for Specific Weeds:

1. Partially controlled.
2. Control of moderate to heavy infestations of nightshade will be improved with a tank mixture of the appropriate labeled rate of an atrazine premix product or a surface applied acetanilide product such as Dual® II herbicide, Dual II Magnum® herbicide, Surpass® herbicide or Topnotch® herbicide.
3. Burndown control of Canada thistle in minimum and no-till corn only.
4. To aid in control of waterhemp, apply Hornet WDG in tank mix combination with the appropriate labeled rate of a surface applied acetanilide product such as Surpass or TopNotch herbicide.
5. Hornet WDG will not control ALS resistant or tolerant biotypes of kochia.

Hornet WDG may be soil applied as a preplant surface, preplant incorporated, or preemergence treatment. Apply alone or in tank mix combination with a grass control product such as Surpass or Topnotch herbicide.

Tank Mixing Limitations: Hornet WDG may be applied in tank mix combination with other products provided (1) the timing and method of application is the same as recommended for Hornet WDG; and (2) tank mixing with Hornet WDG is not prohibited by the label of the tank mix product. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels.

Soil Application Directions

Applications may be made from 30 days prior to planting through V6 stage or 20" tall corn, whichever occurs first.

1. **Preplant Incorporated Application:** For best results, apply and incorporate Hornet WDG from 0 to 30 days before planting. Preplant incorporated treatments may be applied in water or liquid fertilizer. Uniformly incorporate the herbicide treatment into the top 2 to 3 inches of the final seedbed.

2. **Preplant Surface Applied:** For best results in minimum-tillage or no tillage systems, Hornet WDG alone and with certain tank mixtures may be applied up to 30 days before planting. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide such as Gramoxone Extra herbicide, Touchdown herbicide, glyphosate (Glyphomax® Plus herbicide or Roundup UltraMAX herbicide. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. To the extent possible do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Canada Thistle Control in Minimum and No-Till Corn:

Hornet WDG may be applied as a burndown treatment for control of emerged Canada thistle in minimum and no-till corn. The application will result in reduced late season competition. Delay the application until most of the thistle has emerged and averages 4 to 8 inches in height. For applications to Canada thistle, always include crop oil concentrate (See “Adjuvant Systems” in “Postemergence Treatments” section). Tank mix Hornet WDG with glyphosate (Glyphomax Plus or Roundup UltraMAX), or sulfosate (Touchdown) and non-ionic surfactant for burndown control of existing grass and annual broadleaf weeds. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. Do not apply in tank mixture with Gramoxone Extra as this will result in reduced control of Canada thistle. Do not cultivate for at least 14 days after application to allow for thorough translocation of the herbicide treatment.

Note: Hornet WDG will not control Canada thistle that has not emerged at the time of application in minimum or conventional tillage systems.

3. **Burndown Application:** When used as a burndown application, Hornet WDG will provide foliar control of broadleaf weeds listed in the “Postemergence Treatments” section of this label and residual control of weeds listed under soil application. Foliar burndown applications should always include crop oil concentrate (see “Adjuvant Systems” in “Postemergence Treatments” section). To broaden the spectrum of weeds controlled, Hornet WDG may be tank mixed with other herbicides such as glyphosate (Glyphomax Plus or Roundup UltraMAX), sulfosate (Touchdown), paraquat (Gramoxone Extra), or 2,4-D herbicide, etc. (See tank mixing instructions.)
4. **Preemergence Application:** Apply at the time of planting or after planting, but prior to crop or weed emergence. Adequate soil moisture following application is required for optimum herbicidal activity. For surface applications, rainfall, or overhead sprinkler irrigation is necessary to move Hornet WDG into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture, and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is generally adequate. If adequate soil moisture is not received within 7 to 10 days after a surface applied treatment, a shallow cultivation is recommended to control established weeds and move the herbicide into the weed germination zone. When adequate soil moisture is received following dry conditions, performance may vary with weed species and the depth of the weed root system in the soil.

5. **Spike Stage Application:** Apply from corn emergence (ground cracking stage) until corn is 2 inches in height and before the first leaf is unfurled. Adequate soil moisture is required for optimum herbicidal activity. For those weeds that have not emerged at the time of application, rainfall or overhead sprinkler irrigation is necessary to move Hornet WDG into the weed germination zone. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture, and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is generally adequate. If adequate soil moisture is not received within 7 to 10 days after a surface applied treatment, a shallow cultivation is recommended to control established weeds and mix the herbicide into the weed germination zone. When adequate soil moisture is received following dry conditions, performance may vary with weed species and rooting depth of target weeds.

**Tank Mixing
(Preplant Surface Applied, Preplant Incorporated, and Postplant Preemergence Treatments)**

Note: When tank mixing with a companion herbicide, read and follow each manufacturer’s label for weeds controlled, applicable use directions, precautions, and limitations.

1. Reduced Rates of Hornet WDG Plus Atrazine-Containing Pre-Mix Products

Reduced rates of Hornet WDG can be tank mixed with labeled rates of atrazine-containing pre-mix herbicide products such as Surpass 100 herbicide, FulTime® herbicide, Keystone® herbicide or Keystone LA herbicide for improved control of certain broadleaf weeds not consistently controlled by atrazine pre-mix products. Hornet WDG may be applied in tank mix combination with other products provided (1) the timing and method of application is the same as recommended for Hornet WDG; and (2) tank mixing with Hornet WDG is not prohibited by the label of the tank mix product. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. Reduced rates of Hornet WDG tank mixed with labeled rates of these atrazine pre-mix products will provide consistent preemergence control of velvetleaf, lambsquarters, pigweed species, waterhemp, and triazine “resistant” varieties (triazine tolerant biotypes) of these species. These tank mixtures will also provide improved control of large-seeded broadleaf weeds such as cocklebur, common ragweed, giant ragweed, common sunflower, and jimsonweed.

On soils with less than 3% organic matter, tank mix Hornet WDG at 3.0 oz/A with the recommended label rate of the atrazine pre-mix product. On soils with greater than 3% organic matter, tank mix Hornet WDG at 4.0 oz/A with the recommended label rate of the atrazine pre-mix product.

Soil Organic Matter	Hornet WDG (oz/acre)	Acres per 6 lb Plastic Jug	Acres per 6 oz Packet (Contains four 1.5 oz units)
<3%	3.0	32	2
>3%	4.0	24	1.5

2. Hornet WDG plus Glyphosate (Glyphomax Plus or Roundup UltraMAX), Gramoxone Extra, or Touchdown for Minimum-tillage or No-tillage Systems

In minimum-tillage or no-tillage situations where corn is planted directly into a cover crop, stale seedbed, or previous crop residues, herbicides such as glyphosate (Glyphomax Plus or Roundup UltraMAX), paraquat (Gramoxone Extra), or sulfosate (Touchdown) may be tank mixed with Hornet WDG. Apply in 10 to 60 gallons of water or fluid fertilizer per acre with ground equipment. The higher end of the carrier rate will provide better coverage under high residue situations. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels.

Application Timing: Apply before, during (behind the planter), or after planting, but before the crop emerges.

Glyphomax Plus or Roundup UltraMAX: See the product label for Glyphomax Plus, Roundup UltraMAX, (or other labeled glyphosate) herbicide label for weeds controlled, recommended rates for specific weeds, and application instructions.

Gramoxone Extra: See the label for Gramoxone Extra for weeds controlled, recommended rates for specific weeds, and application instructions. Do not apply combinations containing Gramoxone Extra in suspension type fertilizers as the activity of the active ingredient paraquat will be reduced.

3. Hornet Plus 2,4-D for Minimum-tillage or No-tillage Systems

Where heavy crop residues exist, add 1.0 to 2.0 pints per acre of an appropriately labeled 3.8 – 4.0 lb a.e. per gallon 2,4-D amine or ester to the spray tank and apply in a volume of carrier capable of providing sufficient coverage of the crop residue. A carrier volume of 20 gallons per acre is recommended in heavy crop residue situations.

As carriers, nitrogen solutions and complete liquid fertilizers applied before corn emergence will enhance burndown of existing weeds and, therefore, are recommended instead of water. Add a crop oil concentrate or non-ionic surfactant at 1.0 to 2.0 quarts per 100 gallons diluted spray or another appropriate surfactant at its recommended rate. Apply before weeds reach 6 inches in height. This tank mixture will not control emerged grasses.

Hornet WDG Soil-Applied Followed by Postemergence Treatments:

Broadleaf weeds not controlled by Hornet WDG may be controlled with a postemergence herbicide product such as Hornet WDG, Banvel herbicide, Clarity herbicide, 2,4-D herbicide, Marksman herbicide, Buctril herbicide, or Beacon herbicide. Read and follow each manufacturer’s label for weeds controlled, applicable use directions, precautions, and limitations before use.

Postemergence Treatments

Apply Hornet WDG as a postemergence spray at a rate of 2.0 - 5.0 ounces per acre. Use higher rates for control of heavy weed infestations, larger weeds, or when a longer period of residual control is desired. When applied postemergence, Hornet WDG must be used with one of the adjuvant systems described below.

Postemergence Application Rates:

Acres Per Package Type	Application Rate (oz/acre) †			
	2.0 oz/acre	3.0 oz/acre	4.0 oz/acre	5.0 oz/acre
Acres per 6 oz Packet (Contains four 1.5 oz units)	3	2	1.5	1.2
Acres per 6 lb jug	48	32	24	19.2

† Refer to Mixing Directions section to determine the number of water soluble packets and total spray volume required for treated acreage.

Application Timing

Apply to actively growing weeds as a broadcast, or band treatment from the time of corn emergence (spike stage) until corn reaches 20 inches in height or the V6 stage whichever occurs first. For optimal control, apply before broadleaf weeds exceed the maximum height listed. Weeds that exceed the maximum height listed may be suppressed and recover after 2 to 3 weeks.

Directed Postemergence Application: Hornet WDG may be applied as a directed postemergence application to corn that is 20 to 36 inches in height or has more than 6 leaf collars. Use only drop nozzles and avoid spraying the corn plant by directing the spray as low as possible while allowing for optimal coverage of weeds. Use the highest labeled rates for weeds greater than the maximum size listed on this label. Control of weeds larger than the maximum height listed may vary due to weeds species, stage of growth, and growing conditions. Results may range from complete control to suppression.

- Do not spray into the whorl of corn plants.
 - Do not apply to corn more than 36 inches tall.
- Note:** If field corn is grown for forage or ensilage, application must occur before corn reaches 20 inches in height or V6 growth stage (whichever occurs first) and an interval of at least 45 days is required between application and harvest.

Factors Affecting Weed Control: Apply to actively growing weeds. Extreme growing conditions such as drought, or near freezing temperatures before, at, or following application may result in reduced weed control. Degree of control will depend on coverage of treated weeds and weed susceptibility as well as growing conditions at the time of treatment.

Environmental Conditions and Herbicidal Activity of Hornet WDG:

Factors in effective weed control with Hornet WDG include application rate, weed size, daytime temperature, soil moisture prior to and following application, and use of adjuvants. Best weed control results are obtained when Hornet WDG is applied to small, actively growing weeds, when daytime temperatures are warm (70°F or more), and soil moisture is adequate to support active weed growth prior to and following application. If weeds are under drought stress, consider delaying application until more favorable conditions resume. Application when weeds are moisture stressed or taller than the recommended height for control may result in only partial control.

- Hornet WDG is rainfast in 2 hours.
- Applications made immediately prior to, during, or immediately following periods of large day/night temperature fluctuations or where daytime temperatures do not exceed 60°F may decrease weed control.
- Poor weed control may result from applications made to plants under stress from:
 - ▶ abnormally hot or cold weather
 - ▶ environmental conditions such as drought, water-saturated soils, hail damage, or frost
 - ▶ prior herbicide applications

Use of Adjuvants: All postemergence applications of Hornet WDG must include 1) a non-ionic surfactant at 0.25% volume/volume (1 qt/100 gal) or 2) crop oil concentrate or methylated seed oil at 1% volume/volume (1 gal/100 gal). Use a good quality surfactant with at least 80% active ingredient (of which at least 50% is actual non-ionic surfactant). Under extremely dry growing conditions, the use of an agriculturally approved sprayable liquid fertilizer or ammonium sulfate, in combination with the non-ionic surfactant or crop oil concentrate or methylated seed oil may enhance control. Use 28%, 30%, or 32% urea ammonium nitrate at 2.5% volume/volume (2.5 gal/100 gal) or 2 to 4 lb of sprayable grade ammonium sulfate per acre.

Note: Do not use liquid fertilizer solutions or suspensions as the total carrier because excessive crop injury may occur. Use only EPA approved surfactants for use on food crops.

Cultivation: For best results, do not cultivate within 10 days before or after application.

Tank Mixing: Hornet WDG may be applied in tank mix combination with other products provided (1) the timing and method of application is the same as recommended for Hornet WDG; and (2) tank mixing with Hornet WDG is not prohibited by the label of the tank mix product; and (3) the tank mix combination is compatible as determined by a "jar test" described in the "Tank Mix Compatibility Testing" section. When tank mixing, do not exceed recommended application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. For control of grass weeds, Hornet WDG may be tank mixed with a postemergence grass herbicide such as Accent or Basis Gold, Option or Steadfast. For an expanded spectrum of broadleaf weed control, Hornet WDG may be tank mixed with other postemergence broadleaf herbicides such as atrazine, Banvel, Buctril, Callisto, Clarity, Distinct, or 2,4-D. Hornet WDG may also be tank mixed with Glyphomax Plus, Roundup UltraMAX, or other labeled glyphosate formulations for application to Roundup Ready field corn.

Do not post apply Hornet WDG in tank mix combination with Basagran, Laddock or Lightning herbicides as severe crop injury may result.

Weeds Controlled And Application Rates for Postemergence Application

(Use higher rates for control of larger weeds and for control of heavy weed infestations.)

Hornet WDG will control triazine tolerant biotypes of these weeds, commonly know as “triazine resistant”.

Note: Numbers in parentheses (-) within table refer to Specific Use Directions below.

Annual Weed Control			
Application to “Spike” Corn (1)	Postemergence Application After “Spike” Stage of Growth		
4.0 to 5.0 oz/acre	2.0 oz/acre (weeds 1 - 3 in. tall)	3.0 oz/acre (weeds 1 - 6 in. tall)	4.0 oz/acre (weeds 1 - 8 in. tall)
anoda, spurred beggarweed, Florida buckwheat, wild carpetweed chickweed, common cocklebur, common henbit horseweed (marestail) jimsonweed kochia (2) ladysthumb lambsquarters, common mallow, venice mustard, wild nightshade species pigweed, redroot pigweed, smooth poinsettia, wild puncturevine purslane, common ragweed, common shepherd’s purse sicklepod sida, prickly smartweed, Pennsylvania spurge, nodding spurge, prostrate spurge, spotted sunflower, common thistle, Russian velvetleaf waterhemp species	anoda, spurred beggarweed, Florida chickweed, common cocklebur, common henbit horseweed (marestail) mallow, venice mustard, wild poinsettia, wild puncturevine purslane, common shepherd’s purse sida, prickly spurge, nodding spurge, prostrate spurge, spotted sunflower, common velvetleaf	anoda, spurred beggarweed, Florida chickweed, common cocklebur, common henbit horseweed (marestail) jimsonweed ladysthumb mallow, venice mustard, wild poinsettia, wild puncturevine purslane, common ragweed, common ragweed, giant shepherd’s purse sida, prickly smartweed, Pennsylvania spurge, nodding spurge, prostrate spurge, spotted sunflower, common velvetleaf	anoda, spurred beggarweed, Florida chickweed, common cocklebur, common henbit horseweed (marestail) jimsonweed ladysthumb lettuce, prickly mallow, Venice mustard, wild poinsettia, wild puncturevine purslane, common ragweed, common ragweed, giant shepherd’s purse sida, prickly smartweed, Pennsylvania spurge, nodding spurge, prostrate spurge, spotted sunflower, common velvetleaf
Partial Control	Partial Control	Partial Control (weeds <2 in. tall)	Partial Control (weeds <4 in. tall)
morningglory, entireleaf morningglory, ivyleaf morningglory, tall ragweed, giant	jimsonweed ladysthumb ragweed, common ragweed, giant smartweed, Pennsylvania	buckwheat, wild kochia (2) lambsquarters, common lettuce, prickly morningglory, entireleaf morningglory, ivyleaf morningglory, tall nightshade species pigweed, redroot pigweed, smooth sicklepod thistle, Russian waterhemp species (2)	buckwheat, wild kochia (2) lambsquarters, common morningglory, entireleaf morningglory, ivyleaf morningglory, tall nightshade species pigweed, redroot pigweed, smooth sicklepod thistle, Russian waterhemp species (2)

Biennial and Perennial Weed Control			
Apply 3.0 to 5.0 oz/acre to weeds 3 - 9 inches tall (3,4)			
alfalfa, volunteer	clover, red	dock, curly	thistle, Canada (5)
artichoke, Jerusalem	clover, sweet	sorrel, red	wormwood, biennial
burdock, common	dandelion		

Specific Use Directions:

- Spike corn:** Apply 5 oz per acre for greater residual control on soils with greater than 3% organic matter. Apply 5 oz per acre to increase the degree of partial control for morningglory species and giant ragweed.
- Hornet WDG will not control ALS resistant or tolerant biotypes.
- Biennial and Perennial weeds:** A rate of 4.0 - 5.0 oz per acre will generally provide season-long control. A rate of 3.0 oz per acre will provide control of top growth only. Some regrowth may occur by the end of the season.
- Biennial and Perennial weeds:** Do not tank mix with contact herbicides (such as atrazine, metribuzin, or bromoxynil) as reduced weed control will result.
- Canada thistle:** For Canada thistle control the following season, expressed as stand reduction, apply 5.0 oz per acre of Hornet WDG in tank mix combination with 4.0 oz per acre of Stinger herbicide †.

† **Note:** Maximum Use Rate for the active ingredient clopyralid is 0.25 lb per acre. One ounce of Hornet WDG contains 0.031 lb of clopyralid. One fluid ounce of Stinger contains 0.023 lb of clopyralid.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- Refund of purchase price paid by buyer or user for product bought, or
- Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®Trademark of Dow AgroSciences LLC

®Dual MAGNUM and Dual II MAGNUM are registered trademarks of Syngenta

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Label Code: D02-099-009
Replaces Label: D02-099-008
LOES Number: 010-00082

EPA accepted 02/20/2003

Revisions:

- Updated Storage and Disposal